

Short Communication

Differential Distribution of PDE4B Splice Variant mRNAs in Rat Brain and the Effects of Systemic Administration of LPS in Their Expression

ELISABET REYES-IRISARRI,¹ SILVIA PÉREZ-TORRES,¹ XAVIER MIRÓ,² EMILI MARTÍNEZ,¹ PERE PUIGDOMÈNECH,² JOSÉ M. PALACIOS,³ AND GUADALUPE MENGOD^{1*}

¹Department of Neurochemistry, Institut d'Investigacions Biomèdiques de Barcelona, CSIC (IDIBAPS), Barcelona, Spain

²Department of Molecular Genetics, Institut de Biologia Molecular de Barcelona, CSIC, Barcelona, Spain

³Parc Científic, Universitat de Barcelona, Barcelona, Spain

KEY WORDS cAMP; inflammation; PDE4B splicing variant; in situ hybridization

ABSTRACT Phosphodiesterases (PDE) control intracellular cyclic adenosine monophosphate (cAMP) levels, which appear to play an important role in the regulation of inflammation. PDE4B is especially important in this process. Using in situ hybridization histochemistry we first mapped the expression sites of the four PDE4B splicing forms in rat brain. Using the systemic administration of the bacterial endotoxin lipopolysaccharide (LPS) as an inflammation model in rats, we found an increase in PDEB2 mRNA expression in choroid plexus. The differential expression of PDE4B spliced forms and the differential regulation of PDE4B2 in an inflammatory model further supports an involvement of this splicing variant in the inflammatory response. **Synapse** 62:74–79, 2008.

© 2007 Wiley-Liss, Inc.

INTRODUCTION

The phosphodiesterase 4 (PDE4) family, which specifically hydrolyzes intracellular cAMP, is composed by four isozymes (PDE4A, PDE4B, PDE4C, and PDE4D) encoded by different gene loci, and each of them produces several mRNAs by alternative splicing (Houslay et al., 1998). These isozymes and their splice variants are widely expressed in many tissues including the brain (D'Sa et al., 2005; McPhee et al., 2001; Miró et al., 2002; Pérez-Torres et al., 2000). The different knock out mice for PDE4B and PDE4D genes present unique phenotypes, (Jin and Conti, 2002; Jin et al., 1999, 2005), which suggests different functions for these isozymes.

The majority of studies into PDE4 subtype distribution have focused upon inflammatory cells (Engels et al., 1994; Gantner et al., 1998; Uhlig et al., 1997). PDE4A, PDE4B, and PDE4D, are found to some extent in different inflammatory cell types, where they could be important regulators of inflammatory processes. Selective inhibitors of PDE4s have been suggested as therapies for the treatment of several human diseases (Menniti et al., 2006), predominantly disorders of the immune and inflammatory systems (Teixeira et al.,

1997) or disorders of the central nervous system, such as, depression (O'Donnell and Zhang, 2004) or Alzheimer's disease (McGeer and McGeer, 1995).

The lipopolysaccharide (LPS) is a potent inflammatory agent that has been used to characterize the acute inflammatory process in the brain (Breder and Saper, 1996; Breder et al., 1994). The efficacy of PDE4 inhibitors in inflammation models is remarkable and includes inhibition of LPS-induced increment in serum levels of tumor necrosis factor- α (TNF- α) in liver injury, lung injury, renal failure, mortality (Jin and Conti, 2002; Teixeira et al., 1997), or in a multiple sclerosis mouse model (Moore et al., 2006).

The present study aims to determine the regional distribution in rat brain of the four PDE4B splice variants. The possible involvement of cAMP through

Contract grant sponsor: Fundació La Marató de TV3; Contract grant number: 1017/97; Contract grant sponsor: CICYT; Contract grant number: SAF2003-02083, SAF2006-10243

*Correspondence to: Guadalupe Mengod, Department of Neurochemistry, Institut d'Investigacions Biomèdiques de Barcelona, CSIC (IDIBAPS), Rosselló, 161, 6^a, 08036 Barcelona, Spain. E-mail: gmlnqr@iibb.csic.es

Received 26 April 2007; Accepted 17 July 2007

DOI 10.1002/syn.20459

Published online in Wiley InterScience (www.interscience.wiley.com).

PDE4B in inflammation prompted us to analyze the expression of the different PDE4B splicing variants in the brain of LPS treated rats.

MATERIALS AND METHODS

Animal procedure was performed according to the European Union regulations (O.J. of E.C. L358/1 18/12/1986) and was approved by the Institutional Animal Care and Use Committee. Adult male Wistar rats (200–300 g; Iffa Credo, Lyon, France) used for the neuroanatomical localization of PDE4B splicing variants expression were decapitated and the brain was removed, frozen on dry ice, and kept at -20°C . LPS treatment was performed as follows: rats were injected intraperitoneally with vehicle or LPS (from *Escherichia coli*, serotype 055:B5, Sigma, St. Louis, MO; 500 $\mu\text{g}/\text{kg}$ of body weight) diluted in 300 μl of 0.9% sterile saline. Four rats were sacrificed at each of 1, 2, 3, and 4 h postinjection. Eight control rats received an injection of 0.9% sterile saline, and were sacrificed four at each of 2 and 3 h postinjection.

Several oligonucleotides were synthesized for each splicing variant: PDE4B1, bp 383–427, and 506–550, AF202732; PDE4B2 bp 418–462, 520–564, 545–589, L27058; PDE4B3 bp 476–520, 556–600, 616–660, 700–744, U95748; PDE4B4 bp 171–215, 216–260, 264–308, AF202733. COX-2 mRNA oligonucleotides were complementary to bp 1848–1893, NM_017232. The mRNA regions for each PDE4B splice variants were chosen because they share no similarity with each other. Evaluation of the oligonucleotide sequences with basic local alignment search tool of EMBL and GenBank databases indicated that the probes do not present any significant similarity with mRNAs other than their corresponding targets in the rat. The specificity of the autoradiographic signal obtained in the in situ hybridization histochemistry experiments was confirmed by performing a series of routine controls (Pompeiano et al., 1992) such as the use of different oligonucleotides for the same mRNA obtaining identical hybridization patterns with each individual probe in the entire rat brain; competition with the same unlabeled oligonucleotide for the nonspecific hybridization signal, determination of the T_m of the hybrids, etc. The procedures followed for the in situ hybridization experiments have been published elsewhere (Reyes-Irisarri et al., 2005). The results presented in this work have been done using only one of the different oligonucleotides designed for each splicing form appropriately labeled at their 3' end with ^{33}P -dATP.

A semiquantitative measure of the optical densities of the autoradiograms was conducted with the MCID4 and AIS^R computerized image analysis systems (Imaging Research, St Catharines, Ontario, Canada). Anatomical brain structures were verified by the examination of cresyl violet-stained sections and identified using a rat

brain atlas (Paxinos and Watson, 1998). For each rat, individual values of optical densities in each region were calculated as the mean of two adjacent sections in both hemispheres (average of four measures per rat and region). The following regions were measured: area postrema, CA fields of the hippocampus, cerebellum, choroid plexus, corpus callosum, dorsal cochlear nucleus, dentate gyrus, hilus, leptomeninges, medial genicular nucleus, medial habenular nucleus, median eminence, organum vasculosum of the lamina terminalis, primary somatosensory cortex, and pyramidal tract. Statistical analysis was performed using GraphPad Prism 4 (GraphPad Software, San Diego, CA) and SPSS v14 (SPSS, Chicago, IL). Analysis of data variance for all the experiments were performed considering the factors PDE (four levels), brain region (13 levels), and treatment (five levels: control and 1, 2, 3, and 4 h postinjection). Then, the different regions were analyzed by factors PDE and treatment. Post hoc analysis (Bonferroni's test) was performed for treatment by each PDE in the statistically affected regions. Since no significant statistical difference between the two groups of control rats was observed in any of the regions analyzed, we unified the values as a single control group.

RESULTS AND DISCUSSION

mRNA distribution of PDE4B splicing variants in rat brain

We used in situ hybridization histochemistry to study the distribution of the four PDE4B splicing variants at various coronal levels of the rat brain (Fig. 1). Table I summarizes our semiquantitative measurements of the content of each mRNA in different brain regions. The splicing variants identified by hybridization with only one oligonucleotide each (and always the same) show a coincident hybridization pattern in several brain areas, such as granular layer of the cerebellum and piriform cortex. In contrast they are differentially expressed in other brain areas, indicating a possible nonredundant role in the regulation of intracellular cAMP levels.

Interestingly, PDE4B1, PDE4B2, and PDE4B3 are detected in the area postrema whereas PDE4B4 is absent (or expressed at very low levels). This area has been implicated as a chemoreceptor trigger zone for emesis (Borinson and Wang, 1953; Carpenter et al., 1988). The emetic side effects of rolipram are well known, when used as an antidepressant, (Scott et al., 1991). So too are those that other PDE4 inhibitors (such as roflumilast and cilomilast) produce in patients treated with them for asthma or chronic obstructive pulmonary disease (Spina, 2003; Chung, 2006). The fact that we found no expression of PDE4B4 mRNA in the rat area postrema (even though rodents do not vomit) and that it has been proposed that in humans PDE4B4 exon is unlikely to encode a protein (Shepherd




Fig. 1. Regional distribution of PDE4B splice variant mRNAs in rat brain. Film autoradiograms from rat rostral to caudal sections are presented, showing the hybridization pattern of PDE4B1 (A₁–A₆), PDE4B2 (B₁–B₆), PDE4B3 (C₁–C₆), and PDE4B4 (D₁–D₆). Note the strong hybridization signal of PDE4B3 in white matter, and the exclusive PDE4B2 expression in dentate gyrus of the hippocampus. aca, anterior commissure, anterior part; AcbSh, accumbens nucleus, shell; aci, anterior commissure, intrabulbar part; AO, anterior olfactory nucleus; AP, area postrema; Arc, arcuate nucleus; CA, Ammon's

horn of hippocampus; chp, choroid plexus; cp, cerebral peduncle; CPu, caudate putamen (striatum); DCo, dorsal cochlear nucleus; DG, dentate gyrus; gcc, genu of the corpus callosum; hil, hilus; LRt, lateral reticular nucleus; ME, median eminence; MG: medial geniculate nucleus; MHb, medial habenular nucleus; 3, oculomotor nucleus; Pir, piriform cortex; Pn, pontine nucleus; PV, paraventricular thalamic nucleus; py, pyramidal tract; S1: primary somatosensory cortex. Bar = 3 mm.

et al., 2003), points to the importance of determining the expression of PDE4 splice variants in this area and other related areas of the human brain. This will help to find candidate targets for specific splicing variant PDE4 inhibitors with no emetic side effects to treat the aforementioned diseases.

In the hippocampal formation all four isoforms are visualized in CA2 and CA3 fields. In the dentate gyrus the only splicing isoform found was PDE4B2. The basic circuitry of the hippocampus is a trisynaptic circuit (Brown and Zador, 1990) where the sensory inputs from the entorhinal cortex arrive at the gran-

TABLE I. Estimated densities of PDE4B splicing variable mRNAs in different regions of the rat brain

Brain area	PDE4B1 ^a	PDE4B2 ^a	PDE4B3 ^a	PDE4B4 ^a
Cortex				
Parietal cortex	+/++	+	+	+/++
Frontal cortex	+/+	+	+	+/+
Cingulate cortex	+/+	+	+	+
Retrosplenial cortex	+/+	+	+	+
Entorhinal cortex	+/++	+	+	+
Olfactory system				
Anterior olfactory nucleus	-/+	+/++	+/++	+/++
Olfactory tubercle	+	+/+	+/++	+
Piriform cortex	+/+/++	++	++	++
Islands of Calleja	+	+	+	+
Islands of Calleja, major island	+	-/+	+	-/+
Basal ganglia and related areas				
Caudate-Putamen	+/++	-/+	+	-/+
Accumbens				
Core	+	-/+	+/+	-/+
Shell	++	+	+	+
Limbic areas				
Ammon's horn				
CA1 (pyramidal cell layer)	+/+/++	++	+	+
CA2 (pyramidal cell layer)	+/++	++	++	++
CA3 (pyramidal cell layer)	+/++	++	++	+/+
Dentate gyrus	++	+/+/++	-/+	-
Hilus	+	+	+/++	-/+
Subiculum	+	+	+	-/+
Pre, parasubiculum	+	+	+	-/+
Amygdala	+/++	+	+	+
Lateral septal nucleus	+/++	-	-	-/+
Medial septal nucleus	-/+	-/+	+	+
Thalamus and Hypothalamus				
Medial habenular nucleus	-/+	+/++	++	+/++
Lateral habenular nucleus	-/+	+	++	+/++
Paraventricular thalamic nucleus	-	-	++	+/++
Ventroposterior thalamic nuclei	-	+	+	-
Laterodorsal thalamic nucleus	-	+	+	-
Mediodorsal thalamic nucleus	-	+	+	-
Posterior thalamic nucleus group	-	+	+	-
Zona incerta	+	-/+	+/++	-/+
Reticular thalamic nucleus	-	-/+	+	-
Dorsomedial hypothalamus	+/++	-/+	-/+	-
Ventromedial hypothalamus	+/++	+	-/+	-
Arcuate nucleus	-	+/+/++	-	-
Periventricular hypothalamic nucleus	-	+	-	-
Medial geniculate nucleus	-/+	+	+/++	+/++
Internal capsule	-	+	+/+/++	++
Brainstem				
Superior colliculus	+	+	+/++	+
Oculomotor nucleus	-	+	+/++	-/+
Pontine nucleus	++	++	+/++	+/++
Accessory facial nucleus	-	-/+	+	-/+
Facial nucleus	-/+	-/+	+	+
Dorsal cochlear nucleus	+/++	+	++	+
Nucleus of the solitary tract	+	-/+	-	-/+
Hypoglossal nucleus	-/+	-/+	+	-/+
Cuneate nucleus	-/+	-/+	+/++	-/+
Dorsal motor nucleus of vagus	-/+	-/+	-	-/+
Lateral reticular nucleus	-/+	-/+	+/++	-/+
Gracile nucleus	-	-/+	+/++	-
Inferior olive	-	+	+/+/++	+
Cerebellum				
Molecular layer	-	-	-	-
Granular layer	+++	+++	+++	+++
White matter	-	-/+	++	-/+
Cerebellar nuclei	-/+	-/+	+	-
Circumventricular organs				
Choroid plexus	-	-/+	-	-
Area postrema	++	+/++	+/++	-
White matter	-	-	+/+/++	+

^amRNA data are expressed as semiquantitative estimates of hybridization intensity obtained by microdensitometric analysis of film autoradiograms. The levels of hybridization signals are: "+++" very strong; "++" moderate; "+" weak; "-/+" very weak and "-" not detected. "White matter" includes pyramidal tract, genu corpus callosum, cerebral peduncle, and anterior commissure.

ule cells of the dentate gyrus, whose main output—via mossy fiber axons—is the pyramidal neurons of CA3, that send their Schaffer collateral axons to pyramidal neurons of CA1. Some of the synapses of the

hippocampus display a remarkable form of plasticity, long-term potentiation (LTP), which could be relevant to the mnemonic functions of this circuitry (Brown et al., 1988), and is a leading candidate for a synaptic




Fig. 2. Distribution of COX-2 mRNA in rat hippocampus from control (A) and LPS stimulated rats at 2-h postinjection (B). Note the profound transcriptional activation (visualized as a spotlike mRNA signal) detected over blood vessels of the entire brain microvasculature and leptomeninges of LPS-injected rats. Bar = 1 mm.

mechanism for rapid learning in mammals (Brown and Zador, 1990). This form of plasticity has been studied most in the mossy fiber and in the Schaffer collateral axons. Several authors (Barad et al., 1998, 1999; Rose et al., 2005) suggest that rolipram could enhance hippocampus-dependent memory tasks, and reduce cognitive decline associated with neurodegenerative and psychiatric diseases. These studies together with our finding that PDE4B2 is the only splicing variant expressed in the dentate gyrus, suggest that it could be involved in LTP regulation in the mossy fiber axons. The findings of Ahmed and Frey (2003) that PDE4B3 is involved in the regulation of LTP in the hippocampus, do not invalidate our suggestion, since there are different mechanisms for the two LTP systems: the mossy fibers and the Schaffer collateral inputs. PDE4B3 could be involved in the LTP of Schaffer collateral inputs (which take place in the CA3 region where all the splicing variants of PDE4B are expressed), and PDE4B2, could be involved in the LTP system of the mossy fibers output, which originates in the dentate gyrus, where PDE4B2 is the only splicing form expressed.

Systemic administration of LPS: effects on the brain

To study the effects of short-term systemic administration of LPS on the expression of PDE4B splicing variants in rat brain, we measured the corresponding mRNA levels 1, 2, 3, and 4 h after LPS treatment in 15 brain areas (including some circumventricular organs) for the four splicing variant mRNAs. To validate our inflammatory animal model, we analyzed COX-2 mRNA expression in LPS-brains, since COX-2 mRNA is upregulated in endothelial and perivascular cells of rat and mouse brain blood vessels after LPS administration (Cao et al., 1997). As expected, we found that COX-2 mRNA levels were considerably increased 2 h after LPS injection in microvasculature and leptomeninges (Fig. 2). The analysis of the autoradiogram optical densities for the four PDE4B splicing variant mRNAs indicated a strong influence of the factors PDE ($F = 126.750$; $P < 0.0001$) and region ($F = 337.929$; $P < 0.0001$), while the influence of treatment was indicated by the interaction between PDE-treatment ($F = 2.057$; $P < 0.05$). Analysis of var-

iance for factors PDE and treatment at the different regions identifies OVTL, hilus, and choroid plexus as the regions where treatment effects were more consistent. Finally, Bonferroni's test performed for each PDE in these regions showed an exclusively significant increase in mRNA expression of PDE4B2 in choroid plexus when comparing vehicle with 2 and 3 h after LPS injection ($P < 0.01$). In these animal groups mRNA presented a tendency to increase in the rest of the areas.

Upregulation in the expression of several mRNAs has been reported to occur in some of the brain areas we have measured, such as choroid plexus, leptomeninges, and some circumventricular organs (organum vasculosum of the lamina terminalis, median eminence, and area postrema) in inflammatory models. Several authors have observed an overexpression of CD14 (Lacroix et al., 1998), TNF- α (Nadeau and Rivest, 1999; Quan et al., 1999), COX-2 (Breder and Saper, 1996; Lacroix and Rivest, 1998), or IKappaB- α (Quan et al., 1997) mRNAs in these regions after a peripheral injection of LPS. The upregulation of PDE4B2 mRNA we observed in the choroid plexus and circumventricular organs after LPS injection could be of interest, given the fact that they may be the sites where communication between peripheral immune system and the brain takes place. The potential role played by PDE4B in inflammation is supported by the work of Conti's group done in PDE4B knockout mice (Jin and Conti, 2002; Jin et al., 2005). Those animals present a decreased responsiveness to LPS in many inflammatory cells: peripheral leukocytes and peritoneal macrophages are not able to secrete TNF- α after LPS stimulation.

To conclude, we have observed a differential distribution of the four splicing variants of PDE4B in the rat brain and a differential regulation for one of the PDE4B splicing variants, PDE4B2, in an inflammatory model. Additional experiments and the development of specific inhibitors for this isoform will help to elucidate its implication in the inflammatory reaction.

ACKNOWLEDGMENTS

S.P.-T. and X.M. have received a fellowship from CIRIT (Generalitat de Catalunya). The authors wish to acknowledge technical help from Rocío Martín. The authors thank Robin Rycroft for English corrections.

REFERENCES

- Ahmed T, Frey JU. 2003. Expression of the specific type IV phosphodiesterase gene PDE4B3 during different phases of long-term potentiation in single hippocampal slices of rats in vitro. *Neuroscience* 117:627–638.
- Bach ME, Barad M, Son H, Zhuo M, Lu YF, Shih R, Mansuy I, Hawkins RD, Kandel ER. 1999. Age-related defects in spatial memory are correlated with defects in the late phase of hippocampal long-term potentiation in vitro and are attenuated by drugs that enhance the cAMP signaling pathway. *Proc Natl Acad Sci USA* 96:5280–5285.

- Barad M, Bourtchouladze R, Winder DG, Golan H, Kandel E. 1998. Rolipram, a type IV-specific phosphodiesterase inhibitor, facilitates the establishment of long-lasting long-term potentiation and improves memory. *Proc Natl Acad Sci USA* 95:15020–15025.
- Borinson HL, Wang SC. 1953. Physiology and pharmacology of vomiting. *Pharmacol Rev* 5:193–230.
- Breder CD, Saper CB. 1996. Expression of inducible cyclooxygenase mRNA in the mouse brain after systemic administration of bacterial lipopolysaccharide. *Brain Res* 713:64–69.
- Breder CD, Hazuka C, Ghayur T, Klug C, Huginin M, Yasuda K, Teng M, Saper CB. 1994. Regional induction of tumor necrosis factor α expression in the mouse brain after systemic lipopolysaccharide administration. *Proc Natl Acad Sci USA* 91:11393–11397.
- Brown TH, Zador AM. 1990. Hippocampus. In: Gordon MS, editor. *The synaptic organization of the brain*. New York: Oxford University Press. p 346–388.
- Brown TH, Chapman PF, Kairiss EW, Keenan CL. 1988. Long-term synaptic potentiation. *Science* 242:724–728.
- Cao C, Matsumura K, Yamagata K, Watanabe Y. 1997. Involvement of cyclooxygenase-2 in LPS-induced fever and regulation of its mRNA by LPS in the rat brain. *Am J Physiol* 272:R1712–R1725.
- Carpenter DO, Briggs DB, Knox AP, Strominger N. 1988. Excitation of area postrema neurons by transmitters, peptides, and cyclic nucleotides. *J Neurophysiol* 59:358–369.
- Chung KF. 2006. Phosphodiesterase inhibitors in airways disease. *Eur J Pharmacol* 533:110–117.
- D'Sa C, Eisch AJ, Bolger GB, Duman RS. 2005. Differential expression and regulation of the cAMP-selective phosphodiesterase type 4A splice variants in rat brain by chronic antidepressant administration. *Eur J Neurosci* 22:1463–1475.
- Engels P, Fichtel K, Lubbert H. 1994. Expression and regulation of human and rat phosphodiesterase type IV isogenes. *FEBS Lett* 350:291–295.
- Gantner F, Gotz C, Gekeler V, Schudt C, Wendel A, Hatzelmann A. 1998. Phosphodiesterase profile of human B lymphocytes from normal and atopic donors and the effects of PDE inhibition on B cell proliferation. *Br J Pharmacol* 123:1031–1038.
- Houslay MD, Sullivan M, Bolger GB. 1998. The multienzyme PDE4 cyclic adenosine monophosphate-specific phosphodiesterase family: Intracellular targeting, regulation, and selective inhibition by compounds exerting anti-inflammatory and antidepressant actions. *Adv Pharmacol* 44:225–342.
- Jin SL, Conti M. 2002. Induction of the cyclic nucleotide phosphodiesterase PDE4B is essential for LPS-activated TNF- α responses. *Proc Natl Acad Sci USA* 99:7628–7633.
- Jin SL, Richard FJ, Kuo WP, D'Ercole AJ, Conti M. 1999. Impaired growth and fertility of cAMP-specific phosphodiesterase PDE4D-deficient mice. *Proc Natl Acad Sci USA* 96:11998–12003.
- Jin SL, Lan L, Zoudilova M, Conti M. 2005. Specific role of phosphodiesterase 4B in lipopolysaccharide-induced signaling in mouse macrophages. *J Immunol* 175:1523–1531.
- Lacroix S, Rivest S. 1998. Effect of acute systemic inflammatory response and cytokines on the transcription of the genes encoding cyclooxygenase enzymes (COX-1 and COX-2) in the rat brain. *J Neurochem* 70:452–466.
- Lacroix S, Feinstein D, Rivest S. 1998. The bacterial endotoxin lipopolysaccharide has the ability to target the brain in upregulating its membrane CD14 receptor within specific cellular populations. *Brain Pathol* 8:625–640.
- McGeer PL, McGeer EG. 1995. The inflammatory response system of brain: Implications for therapy of Alzheimer and other neurodegenerative diseases. *Brain Res Rev* 21:195–218.
- McPhee I, Cochran S, Houslay MD. 2001. The novel long PDE4A10 cyclic AMP phosphodiesterase shows a pattern of expression within brain that is distinct from the long PDE4A5 and short PDE4A1 isoforms. *Cell Signal* 13:911–918.
- Menniti FS, Faraci WS, Schmidt CJ. 2006. Phosphodiesterases in the CNS: Targets for drug development. *Nat Rev Drug Discov* 5:660–670.
- Miró X, Pérez-Torres S, Puigdomènech P, Palacios JM, Mengod G. 2002. Differential distribution of PDE4D splice variant mRNAs in rat brain suggests association with specific pathways and presynaptic localization. *Synapse* 45:259–269.
- Moore CS, Owens T, Earl N, Frenette R, Styhler A, Nicholson DW, Mancini JA, Hebb AL, Robertson GS. 2006. Peripheral PDE4 inhibition produced by L-826,141 prevents experimental autoimmune encephalomyelitis. *J Pharmacol Exp Ther* 319:63–72.
- Nadeau S, Rivest S. 1999. Regulation of the gene encoding tumor necrosis factor α (TNF- α) in the rat brain and pituitary in response in different models of systemic immune challenge. *J Neuropathol Exp Neurol* 58:61–77.
- O'Donnell JM, Zhang HT. 2004. Antidepressant effects of inhibitors of cAMP phosphodiesterase (PDE4). *Trends Pharmacol Sci* 25:158–163.
- Paxinos G, Watson C. 1998. *The rat brain in stereotaxic coordinates*. San Diego: Academic Press.
- Pérez-Torres S, Miró X, Palacios JM, Cortés R, Puigdomènech P, Mengod G. 2000. Phosphodiesterase type 4 isozymes expression in human brain examined by *in situ* hybridization histochemistry and [3 H]rolipram binding autoradiography. Comparison with monkey and rat brain. *J Chem Neuroanat* 20:349–374.
- Pompeiano M, Palacios JM, Mengod G. 1992. Distribution and cellular localization of mRNA coding for 5-HT $_1$ A receptor in the rat brain: Correlation with receptor binding. *J Neurosci* 12:440–453.
- Quan N, Whiteside M, Kim L, Herkenham M. 1997. Induction of inhibitory factor κ B α mRNA in the central nervous system after peripheral lipopolysaccharide administration: An *in situ* hybridization histochemistry study in the rat. *Proc Natl Acad Sci USA* 94:10985–10990.
- Quan N, Stern EL, Whiteside MB, Herkenham M. 1999. Induction of pro-inflammatory cytokine mRNAs in the brain after peripheral injection of subseptic doses of lipopolysaccharide in the rat. *J Neuroimmunol* 93:72–80.
- Reyes-Irisarri E, Perez-Torres S, Mengod G. 2005. Neuronal expression of cAMP-specific phosphodiesterase 7B mRNA in the rat brain. *Neuroscience* 132:1173–1185.
- Rose GM, Hopper A, De Vivo M, Tehim A. 2005. Phosphodiesterase inhibitors for cognitive enhancement. *Curr Pharm Des* 11:3329–3334.
- Scott AI, Perini AF, Shering PA, Whalley LJ. 1991. In-patient major depression: Is rolipram as effective as amitriptyline? *Eur J Clin Pharmacol* 40:127–129.
- Shepherd M, McSorley T, Olsen AE, Johnston LA, Thomson NC, Baillie GS, Houslay MD, Bolger GB. 2003. Molecular cloning and subcellular distribution of the novel PDE4B4 cAMP-specific phosphodiesterase isoform. *Biochem J* 370:429–438.
- Spina D. 2003. Phosphodiesterase-4 inhibitors in the treatment of inflammatory lung disease. *Drugs* 63:2575–2594.
- Teixeira MM, Gristwood RW, Cooper N, Hellewell PG. 1997. Phosphodiesterase (PDE)4 inhibitors: Anti-inflammatory drugs of the future? *Trends Pharmacol Sci* 18:164–171.
- Uhlig S, Featherstone RL, Held HD, Nusing R, Schudt C, Wendel A. 1997. Attenuation by phosphodiesterase inhibitors of lipopolysaccharide-induced thromboxane release and bronchoconstriction in rat lungs. *J Pharmacol Exp Ther* 283:1453–1459.